arXiv:1302.5024v1 [hep-th] 20 Feb 2013

Closed string Ramond-Ramond proposed higher derivative interactions on fermionic amplitudes in IIB

Ehsan Hatefi

International Centre for Theoretical Physics
Strada Costiera 11, Trieste, Italy [1]

Abstract

The closed form of the correlation functions and the complete form of the amplitude of one closed string Ramond-Ramond (C-field), two fermionic strings and one scalar field in IIB super string theory has been computed in detail. Deriving $< V_C V_{\bar{\psi}} V_\psi V_\phi >$ through using suitable gauge fixing, we discover some new vertices and their infinite higher derivative corrections. We produce both infinite gauge and scalar $u-$channel poles of this amplitude . In particular, by setting the fact that the kinetic term of fermion fields has no correction, employing Born-Infeld action, the Wess-Zumino terms and their higher derivative corrections, we discover all the infinite $t, s-$channel fermion poles. The couplings between one RR and two fermions and all their infinite higher derivative corrections have been explored. In order to look for all the infinite $(s+t+u)-$ channel scalar/gauge poles for $p+2=n, p=n$ cases , we obtain the couplings between two fermions-two scalars and two fermions, one scalar and one gauge field as well as all their infinite higher derivative corrections. Specifically we make various comments based on arXiv:1205.5079 in favor of the universality conjecture for all order higher derivative corrections (with or without low energy expansion) and the relation of open/closed string that is responsible for all super string scattering amplitudes in IIA,IIB .

---

[1] ehatefi@ictp.it

1 Introduction

D-branes [1][2][3] have been clarifying essential tools in most of the progresses in physics and in particular in Super string theories. There are evidences such that D-branes will be the essential setup for deriving Ads/CFT. As an instance of their dynamical aspects, one may talk about the transition between both open and closed string. D-brane physics has diverse dual description (see string dualities [4]). Some more examples like $D0/D4$ system with their explanations have been realized in the introduction of [5]. In particular in [6] we have shown that how the world volume must appear from super gravity 's point of view , namely we set new kind of ADM reduction to IIB which was reduced to H^5 space in 5D and showed that A(ds) brane world might have been understood.

By using the correct boundary conditions [7], it is believed that D$_p$-branes are hypersurfaces in flat spacetime .

In IIA(IIB) string theories, D$_p$-branes with even (odd) p-are related to BPS branes in which there is no instability and supersymmetry is not broken. Apart from these properties, BPS branes do carry Ramond-Ramond charge. To describe dynamics of branes, one has to work out with some proper effective actions, namely one has to deal with bosonic effective actions and also distinguish it from its supersymmetric version. For bosonic actions in the presence of various D$_p$-brane configurations one has to look at the basic reference [8].

To our knowledge the super symmetrized version of that effective action has not been completely found yet, however it is worth giving some important reference [9]. In this paper, in order to avoid some details we would like to highlight and more specifically to just address some of the attempts which have been carried out in these research topics. One might start reading [10] to explore the effective action for a bosonic D$_p$-brane. To follow in detail the derivation of the supersymmetric action once more for a D$_p$-brane one should take in to account [11]. In order to start with a more comprehensive review of Myers terms, the Chern-Simons action (for mixed couplings between closed string RR and scalar fields), the Wess-Zumino actions (mixed couplings between closed string RR and gauge/scalar fields) and more significantly for Born-Infeld action one may want to go through [12, 13] and all their references therein. All the three standard effective field theory methods including Pull-back, Taylor expansion and Myers terms have been realized in [14]. More specifically, to extract the necessary vertices from the effective field theory, the method for obtaining the higher derivative corrections of Myers, Chern-Simons and also Born-Infeld actions, have been extensively expressed in the Section five of [14].

Working out in detail with D_p-brane physics and employing scattering amplitude techniques , one might hope to realize the importance of open strings in which they provide diverse roles/results and probably they would provide fascinating recent physics in near future as well. As an example we may hint a conjecture in [15] in which BPS open strings's quantum effects might indicate namely the host branes's curvature.

In the introduction of [13], we have made several motivations/reasons for our continuous works , nevertheless it is worth pointing out once more the following points.

Given some attempts [16],[17] and [18] it would be nice to find the complete form of WZ and DBI actions which is still incomplete. We provide some more data and our S-matrix will be of help of its determination.

Having crucial tools for higher point functions (especially in string theory scattering amplitudes) is worth for various arguments such as deriving AdS/CFT correspondence. Let us refer to [19] out of so many works as basic papers that involved either the scattering from stable branes or dealt with intriguing applications of the branes.

Within Ads/CFT, there is a concrete relation between a closed and an open string, therefore it is of high importance to actually study the mixed amplitudes, namely discovering open/closed amplitudes with all their higher derivative corrections in detail might provide new light in understanding future works. As an example , the new WZ couplings and their higher derivative corrections (which are found by direct S-matrix computations) have been applied to realize the N^3 entropy of M5 brane, for further details section 4.2 of [5] is suggested.

These corrections will come from the couplings of branes with lower dimension with closed strings meanwhile lower dimensional branes should have been realized as some solitons. To be more specific in [5] we explored some solutions by solving lower dimensional branes within branes with higher dimensions [20]. Another instance is about $D(-1)/D3$ configuration where this system does have N^2 behaviour and this result can be interpreted by considering higher order α' Myers terms. Diverse applications for new couplings with their corrections in M-theory have been discovered [21, 20].

In this paper we work on the complete form of the amplitude of one RR,two fermions and one scalar field in IIB super string theory where both fermions do carry the same chirality. We also find out the closed form of the correlators of $< V_C V_\phi V_{\bar\psi} V_\psi >$.

Having set some of the past works on Myers terms and Wess-Zumino(WZ) effective actions both for BPS [12],[22],[23] and non-BPS branes [24],[25] and [26], we would like to explore all the infinite two fermions-two scalars and also two fermions-one scalar and one gauge field couplings as well as all their infinite higher derivative corrections in such a way that at the end, we might be able to produce all the infinite scalar/ gauge $(t+s+u)$-channel poles of the string theory (for $p+2=n$,$p=n$ where p is spatial dimension of brane and n is related to the rank of H which is field strength of RR). In the other words we do want to discover all the infinite effective couplings between a RR, two fermions, a gauge and one scalar field by matching field theory vertices with an infinite number of gauge/scalar or fermion poles of $< V_C V_\phi V_{\bar\psi} V_\psi >$ of string amplitude. These new couplings which we are getting to derive might have some applications to F-theory as well as M-theory [27] [21, 20],[5].

In order to do so, one should obtain two fermions-two scalars , and two fermions-one gauge-one scalar couplings and all their infinite higher derivative corrections in IIB super string theory.

The vertex of RR, looks like to the fermions's vertices so one may suppose that, the prescription for $< V_C V_A V_A V_A >$ in [12] or $< V_C V_\phi V_A V_A >$ in [22] can be applied here as well, however we will observe the fact that for two fermion-one gauge-one scalar couplings we are not allowed to make use of the kinetic term of the gauge field. Because here there is no external gauge field for $< V_C V_\phi V_{\bar\psi} V_\psi >$ so these kinds of couplings have to be searched just by comparison with the field theory vertices with string theory S-matrix and eventually their coefficients could be fixed in a correct manner. The complete form of two gauge two scalar couplings to all orders in α' were found by comparing with string theory amplitude of $< V_C V_\phi V_A V_A >$ in [22].

This work illustrates the fact that Super Yang-Mills vertices and in particular two fermion-two scalar couplings (also two fermion-one gauge-one scalar couplings) give rise precisely the same scalar/ gauge poles as those were appeared in $< V_C V_\phi V_{\bar\psi} V_\psi >$. In the next publication [28] we show that the same arguments must be held and among the other things we will derive all infinite two fermion-two gauge couplings of $< V_C V_A V_{\bar\psi} V_\psi >$ for various p, n cases.

This paper provides some more information on the universal behaviour of the higher derivative corrections [23] even for fermionic amplitudes. Thus our S-matrix does serve one more test for universal conjecture which is useful in understanding the full DBI,WZ effective actions. It might be also useful for deriving all the singularities of higher point functions of either BPS or non-BPS amplitudes.

This universal behaviour should have origins coming from the deep relation of a closed and one open string. This relation may clarify closed string's composite nature only in terms of open strings. We describe it further in the conclusion section.

In the next section, we explain the string computations of a closed string RR field, two fermion fields and one scalar field in IIB super string theory ($< V_C V_\phi V_{\bar\psi} V_\psi >$).

Then by expanding the amplitude at low energy limit and by finding the desired vertices such as the vertex of one RR,two fermion fields to all orders in α' and also using WZ terms we will produce both infinite scalar and gauge u-channel poles for various cases such as $p+2=n$,$p=n$ cases. We also move on to produce those infinite $t,s-$ channel fermion poles .

Eventually we summarize our results and talk about the deep relation that does exist between a closed string and one open string for the matching of the super string amplitudes. We also mention some proposals for future researches [28]. In order to avoid many details and to get familiar with the notations, Appendix A and B of [14, 22] should be considered.

2 Notations and Analysis of $< V_{\bar\psi} V_\psi V_\phi >$

In the following sections we do want to set CFT methods to actually find out the entire amplitude of a closed string RR (C-field),two fermions (with the same chirality) and one scalar field in the world volume of IIB super string theory.

Given the vast recent research works on scattering amplitudes, it is indeed impossible to address all attempts on this research topic, however one can naively address some of the attempts which have been extensively carried out at least for tree level computations [29, 30, 31, 12, 22, 23]. The needed vertex operators for our purpose are [2]

$$V_\phi^{(0)}(x) \;=\; \xi_i\Big(\partial X^i(x)+\alpha' ik{\cdot}\psi\psi^i(x)\Big)e^{\alpha' ik\cdot X(x)},$$

---

[2] We clearly consider α' in this paper .

$$\begin{aligned}
V_{\phi}^{(-1)}(x) &= e^{-\phi(x)}\xi_i\psi^i(x)e^{\alpha' ik\cdot X(x)}\\
V_{\phi}^{(-2)}(y) &= e^{-2\phi(y)}\xi_i\Big(\partial X^i(y)+\alpha' ik\cdot\psi\psi^i(y)\Big)e^{\alpha' ik\cdot X(y)},\\
V_{\bar\Psi}^{(-1/2)}(x) &= \bar u^A e^{-\phi(x)/2}S_A(x)\,e^{2iq.X(x)}\\
V_{\Psi}^{(-1/2)}(x) &= u^B e^{-\phi(x)/2}S_B(x)\,e^{2iq.X(x)}\\
V_C^{(-\frac{1}{2},-\frac{1}{2})}(z,\bar z) &= (P_-\not{H}_{(n)}M_p)^{\alpha\beta}e^{-\phi(z)/2}S_\alpha(z)e^{i\frac{\alpha'}{2}p\cdot X(z)}e^{-\phi(\bar z)/2}S_\beta(\bar z)e^{i\frac{\alpha'}{2}p\cdot D\cdot X(\bar z)}, \qquad (1)
\end{aligned}$$

(p,q,k) become the momenta of the RR, fermion and scalar field accordingly. Their on-shell condition is $k^2=q^2=p^2=0$. Note that u^A is the fermion 's wave function of Majorana-Weyl in ten dimension of space-time.

Spin indices have been raised by $C^{\alpha\beta}$ which is the charge conjugation matrix,

$$(P_-\not{H}_{(n)})^{\alpha\beta}=C^{\alpha\delta}(P_-\not{H}_{(n)})_\delta{}^\beta \qquad (2)$$

and also the trace may be found as

$$\begin{aligned}
\mathrm{Tr}\,(P_-\not{H}_{(n)}M_p\gamma^k) &\equiv (P_-\not{H}_{(n)}M_p)^{\alpha\beta}(\gamma^kC^{-1})_{\alpha\beta}\\
\mathrm{Tr}\,(P_-\not{H}_{(n)}M_p\Gamma^{iab}) &\equiv (P_-\not{H}_{(n)}M_p)^{\alpha\beta}(\Gamma^{iab}C^{-1})_{\alpha\beta} \qquad (3)
\end{aligned}$$

where P_- is a projection operator, $P_-=\frac{1}{2}(1-\gamma^{11})$, and the field strength of RR becomes

$$\not{H}_{(n)}=\frac{a_n}{n!}H_{\mu_1\ldots\mu_n}\gamma^{\mu_1}\ldots\gamma^{\mu_n}\ ,$$

where for type IIB theory we do have $n=1,3,5$ and $a_n=1$.

Final remark for notation is in order. We set doubling trick to deal with just holomorphic functions (further details should be looked for in Appendix A of [14]).

The amplitude of two fermions and one gauge field has been found out in [32],however, as a matter of warming up and in particular in order to get familiar with notation let us work out with the simplest tree level amplitude of two fermions and one scalar field $<V_{\bar\psi}V_\psi V_\phi>$. This three point function at disk level is given by

$$\mathcal{A}^{\bar\Psi,\Psi,\phi} \sim \sum_{\rm non-cyclic}\int dx_1dx_2dx_3\,\mathrm{Tr}\,\langle V_{\bar\Psi}^{(-1/2)}(x_1)V_{\Psi}^{(-1/2)}(x_2)V_{\phi}^{(-1)}(x_3)\rangle \qquad (4)$$

Substituting the vertex operators, and taking into account the following holomorphic correlators

$$
\begin{aligned}
\langle X^\mu(z)X^\nu(w)\rangle &= -\frac{\alpha'}{2}\eta^{\mu\nu}\log(z-w),\\
\langle \psi^\mu(z)\psi^\nu(w)\rangle &= -\frac{\alpha'}{2}\eta^{\mu\nu}(z-w)^{-1}\ ,\\
\langle \phi(z)\phi(w)\rangle &= -\log(z-w)\ .
\end{aligned}
\tag{5}
$$

(4) can be written down as

$$
\begin{aligned}
\mathcal{A}^{\bar{\psi},\psi,\phi} \sim & \int dx_1dx_2dx_3\xi_{1i}x_{12}^{-1/4}(x_{13}x_{23})^{-1/2}|x_{12}|^{\alpha'^2k_1.k_2}|x_{13}|^{\alpha'^2k_1.k_3}|x_{23}|^{\alpha'^2k_2.k_3}\\
&\times <: S_A(x_1) : S_B(x_2) : \psi^i(x_3) :> \bar{u}_1^A u_2^B
\end{aligned}
\tag{6}
$$

using

$$
<: S_A(x_1) : S_B(x_2) : \psi^i(x_3) :>= 2^{-1/2}x_{12}^{-3/4}(x_{31}x_{32})^{-1/2}(\gamma^i)_{AB}.
$$

and showing the $SL(2,R)$ invariance of the amplitude; we do gauge fixing as $(x_1,x_2,x_3) = (0,1,\infty)$. Setting this gauge fixing into (6), the amplitude becomes

$$
\mathcal{A}^{\bar{\Psi},\Psi,\phi} = 2^{-1/2}\bar{u}_1^A\gamma^i_{AB}u_2^B\xi_i\left(\mathrm{Tr}\left(\lambda_1\lambda_2\lambda_3\right) - \mathrm{Tr}\left(\lambda_1\lambda_3\lambda_2\right)\right) \tag{7}
$$

The final result of string theory can be reproduced in field theory by extracting the kinetic term of the fermion fields $2\pi\alpha'\bar{\psi}\gamma^\mu D_\mu\psi$ where one has to extract the covariant derivative of fermion field and take into account the commutator in its definition $D^i\psi = \partial^i\psi - i[\phi^i,\psi]$.

It is good to know that the Wick-like rule [33, 34] has already been extended in [25, 26, 12, 14] to find the various correlators of two spin operators and an infinite number of fermions and or currents .

2.1 The complete and closed form of $< V_C V_\phi V_{\bar{\psi}} V_\psi >$ amplitude

Having provided the needed details, we are now ready to start finding out the complete form (entire result in all α' orders) of the the scattering amplitude of one closed string RR(in the bulk), two fermions with the same chitrality and a scalar field. Regarding the chirality of fermions our computation makes sense just in IIB super string theory and the entire result can not be extended to IIA (femions do have different chirality in IIA). As a matter of fact neither there is any gauge nor scalar $(t+s+u)-$ channel pole in IIA.

Certainly two fermions two scalars and two fermions,one gauge one scalar corrections which we are going to discover can not be used in IIA. The other point is that we show that field theory must be fine tuned in the sense that there is no any double pole in string theory, even though some double poles might be allowed in field theory (and in particular some Feynman diagrams are allowed to have double poles). Nevertheless we trust on string amplitude. This $< V_C V_\phi V_{\bar\psi} V_\psi >$ amplitude can be looked for as follows

$$\mathcal{A}^{C\phi\bar\psi\psi} \sim \int dx_1dx_2dx_3dzd\bar z\, \langle V_\phi^{(0)}(x_1)V_{\bar\psi}^{(-1/2)}(x_2)V_\psi^{(-1/2)}(x_3)V_{RR}^{(-\frac{1}{2},-\frac{1}{2})}(z,\bar z)\rangle, \qquad (8)$$

We certainly deal with disk level amplitude and one has to embed all 3 open strings on the boundary where RR should be located inside of the disk, also notice that we do clearly want to keep track of $\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3)$ ordering. By replacing all the vertex operators one might reveal that the amplitude should be divided to two different parts $(\mathcal{A}_1,\mathcal{A}_2)$. The final result is complicated so we decided to carry out each part of the amplitude separately. First we look for $(\mathcal{A}_1)$, for this part we need to know the correlation function of four spin operators (with the same chirality) in ten dimensions which can be found in [35] thus we just replace it into the amplitude and the result should be read off as

$$\begin{aligned}\mathcal{A}_1^{C\phi\bar\psi\psi} &\sim \int dx_1dx_2dx_3dx_4dx_5\,(P_-\not{H}_{(n)}M_p)^{\gamma\delta}\xi_{1i}\bar u_1^A u_2^B(x_{23}x_{24}x_{25}x_{34}x_{35}x_{45})^{-1}\\ &\times\frac{1}{2}[(\gamma^\mu C)_{\gamma\delta}(\gamma_\mu C)_{AB}x_{43}x_{52}-(\gamma^\mu C)_{\gamma B}(\gamma_\mu C)_{A\delta}x_{45}x_{23}]I_1\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3), \qquad (9)\end{aligned}$$

where $x_{ij}=x_i-x_j, x_4=z=x+iy, x_5=\bar z=x-iy$, and

$$I_1 = <:\partial X^i(x_1)e^{\alpha' ik_1.X(x_1)}:e^{\alpha' ik_2.X(x_2)}:e^{\alpha' ik_3.X(x_3)}:e^{i\frac{\alpha'}{2}p.X(x_4)}:e^{i\frac{\alpha'}{2}p.D.X(x_5)}:>$$

Using Wick theorem one gets

$$I=\left(\frac{ip^ix_{54}}{x_{14}x_{15}}\right)|x_{12}|^{\alpha'^2k_1.k_2}|x_{13}|^{\alpha'^2k_1.k_3}|x_{14}x_{15}|^{\frac{\alpha'^2}{2}k_1.p}|x_{23}|^{\alpha'^2k_2.k_3}|x_{24}x_{25}|^{\frac{\alpha'^2}{2}k_2.p}|x_{34}x_{35}|^{\frac{\alpha'^2}{2}k_3.p}|x_{45}|^{\frac{\alpha'^2}{4}p.D.p}$$

By replacing I_1, I in the amplitude, we realize that the amplitude is SL(2,R) invariant. Having set the following Mandelstam variables

$$s = \frac{-\alpha'}{2}(k_1+k_3)^2, \quad t=\frac{-\alpha'}{2}(k_1+k_2)^2, \quad u=\frac{-\alpha'}{2}(k_3+k_2)^2$$

and in particular carrying out a special gauge fixing as $(x_1=0,x_2=1,x_3=\infty)$,one can obtain the first part of the amplitude as

$$
\begin{aligned}
\mathcal{A}_1^{C\phi\bar{\psi}\psi} &\sim (P_-H\!\!\!/_{(n)}M_p)^{\gamma\delta}\xi_{1i}\bar{u}_1^A u_2^B(\frac{-ip^i}{2})\int dzd\bar{z}|z|^{2t+2s-2}|1-z|^{2t+2u-2}(z-\bar{z})^{-2(t+s+u)},\\
&\times[(\gamma^\mu C)_{\gamma\delta}(\gamma_\mu C)_{AB}(1-\bar{z})+(z-\bar{z})(\gamma^\mu C)_{\gamma B}(\gamma_\mu C)_{A\delta}]\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3),
\end{aligned}
\tag{10}
$$

These integrals on the closed string position can be done to actually get the entire result (we propose [36] and the Appendix B of [14] for further details), therefore the complete form of the first part of the amplitude is appeared as

$$
\begin{aligned}
\mathcal{A}_1^{C\phi\bar{\psi}\psi} &\sim (P_-H\!\!\!/_{(n)}M_p)^{\gamma\delta}\xi_{1i}\bar{u}_1^A u_2^B(\frac{-ip^i}{2})\\
&\times\Big[(\gamma^\mu C)_{\gamma\delta}(\gamma_\mu C)_{AB}(stL_1+\frac{1}{2}L_2)+(\gamma^\mu C)_{\gamma B}(\gamma_\mu C)_{A\delta}L_2\Big]\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3),
\end{aligned}
\tag{11}
$$

with

$$
\begin{aligned}
L_1 &= (2)^{-2(t+s+u)}\pi\frac{\Gamma(-u)\Gamma(-s)\Gamma(-t)\Gamma(-t-s-u+\frac{1}{2})}{\Gamma(-u-t+1)\Gamma(-t-s+1)\Gamma(-s-u+1)},\\
L_2 &= (2)^{-2(t+s+u)+1}\pi\frac{\Gamma(-u+\frac{1}{2})\Gamma(-s+\frac{1}{2})\Gamma(-t+\frac{1}{2})\Gamma(-t-s-u+1)}{\Gamma(-u-t+1)\Gamma(-t-s+1)\Gamma(-s-u+1)}
\end{aligned}
\tag{12}
$$

This part of the amplitude is not vanished for different cases, for example for $n=p+2$ the first term in (11) has infinite singularities in u-channel and in particular it involves many contact terms. As it is known the expansion is low energy expansion(which reflects the fact that all Mandelstam variables must tend to zero, for further details on the expansions see [12]). Hence it becomes obvious that the first part of the amplitude(just its first term) does include an infinite number of massless poles, however, we postpone its field theory computations to the next section to see which kinds of open strings, namely gauge/ scalar or fermion can be propagated. Let us move to the second part of the amplitude.

Having replaced the vertex operators , it is given by

$$
\begin{aligned}
\mathcal{A}_2^{C\phi\bar{\psi}\psi} &\sim \int dx_1dx_2dx_3dx_4dx_5\,(P_-H\!\!\!/_{(n)}M_p)^{\gamma\delta}\xi_{1i}(2ik_{1a})\bar{u}_1^\alpha u_2^\beta(x_{23}x_{24}x_{25}x_{34}x_{35}x_{45})^{-1/4}\\
&\times<\psi^a\psi^i(x_1):S_\alpha(x_2)::S_\beta(x_3)::S_\gamma(x_4)::S_\delta(x_5):>I\,\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3),
\end{aligned}
\tag{13}
$$

in which

$$
I=|x_{12}|^{\alpha'^2k_1.k_2}|x_{13}|^{\alpha'^2k_1.k_3}|x_{14}x_{15}|^{\frac{\alpha'^2}{2}k_1.p}|x_{23}|^{\alpha'^2k_2.k_3}|x_{24}x_{25}|^{\frac{\alpha'^2}{2}k_2.p}|x_{34}x_{35}|^{\frac{\alpha'^2}{2}k_3.p}|x_{45}|^{\frac{\alpha'^2}{4}p.D.p}
$$

The only subtlety in the second part of the amplitude is how to derive the correlation function between four spin operators (with the same chirality) and one current. Here we summarize the procedure to actually get it as follows.

First we need to take into account the following OPE

$$:\psi^\mu\psi^\nu(x_1):S_\alpha(x_4): \quad \sim \quad -\frac{1}{2}(\Gamma^{\mu\nu})_\alpha^{\beta}S_\beta(x_4)x_{14}^{-1} \tag{14}$$

in which $\Gamma^{\mu\nu}=\frac{1}{2}(\gamma^\mu\gamma^\nu-\gamma^\nu\gamma^\mu)$ and the next step is in fact replacing this OPE (14) into

$$<\psi^a\psi^i(x_1):S_\alpha(x_2)::S_\beta(x_3)::S_\gamma(x_4)::S_\delta(x_5):>$$

and make use of the rest of the correlator which is the correlator of four spin operators (which has been given in [35]). Of course one has to mention the fact that we must apply the same formalism for the other OPE's and finally add them up where the final answer has rather complicated result. One unusual check after having replaced the final answer of that correlator into the amplitude is about the SL(2,R) invariance of the amplitude in which our result does satisfy that condition. The other test of our computation is to produce the leading singularities in which our calculation does pass this test as well.

We gauge fix it as before and we evaluate the integrals on closed string location. Let us write down the final answer for the second part of the amplitude:

$$\mathcal{A}_2^{C\phi\bar\psi\psi} \quad = \quad \Big(\mathcal{A}_{21}+\mathcal{A}_{22}+\mathcal{A}_{23}+\mathcal{A}_{24}+\mathcal{A}_{25}+\mathcal{A}_{26}\Big)\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3)$$

such that

$$\mathcal{A}_{21} \quad \sim \quad \frac{-1}{4}(P_-\not{H}_{(n)}M_p)^{\gamma\delta}\xi_{1i}(2ik_{1a})\bar{u}_1^\alpha u_2^\beta(\Gamma^{ai\mu}C)_{\alpha\beta}(\gamma_\mu C)_{\gamma\delta}\Big[\frac{1}{2}u(s+t)L_1+L_3(-s-t)\Big]$$

$$\mathcal{A}_{22} \quad \sim \quad \frac{-1}{4}(P_-\not{H}_{(n)}M_p)^{\gamma\delta}\xi_{1i}(2ik_{1a})\bar{u}_1^\alpha u_2^\beta(\Gamma^{ai\mu}C)_{\alpha\delta}(\gamma_\mu C)_{\gamma\beta}\Big[L_1(us)-\frac{1}{2}L_2\Big]$$

$$\mathcal{A}_{23} \quad \sim \quad \frac{1}{4}(P_-\not{H}_{(n)}M_p)^{\gamma\delta}\xi_{1i}(2ik_{1a})\bar{u}_1^\alpha u_2^\beta(\Gamma^{ai\mu}C)_{\gamma\beta}(\gamma_\mu C)_{\alpha\delta}\Big[L_1(ut)-\frac{1}{2}L_2\Big]$$

$$\mathcal{A}_{24} \quad \sim \quad \frac{1}{4}(P_-\not{H}_{(n)}M_p)^{\gamma\delta}\xi_{1i}(2ik_{1a})\bar{u}_1^\alpha u_2^\beta(\Gamma^{ai\mu}C)_{\gamma\delta}(\gamma_\mu C)_{\alpha\beta}\Big[L_1(st)+\frac{1}{2}L_2\Big]$$

$$\begin{aligned}\mathcal{A}_{25} &\sim \frac{1}{4}(P_-H\!\!\!/_{(n)}M_p)^{\gamma\delta}\xi_{1i}(2ik_{1a})\bar{u}_1^\alpha u_2^\beta\Big[u(s+t)L_1+L_2\Big]\\ &\quad\times\Big((\gamma^iC)_{\gamma\beta}(\gamma^aC)_{\alpha\delta}-(\gamma^iC)_{\alpha\delta}(\gamma^aC)_{\gamma\beta}\Big)\\[2ex] \mathcal{A}_{26} &\sim \frac{1}{4}(P_-H\!\!\!/_{(n)}M_p)^{\gamma\delta}\xi_{1i}(2ik_{1a})\bar{u}_1^\alpha u_2^\beta\Big[-\frac{1}{2}u(s-t)L_1+(-t+s)L_3\Big]\\ &\quad\times\Big(-(\gamma^iC)_{\gamma\delta}(\gamma^aC)_{\alpha\beta}+(\gamma^iC)_{\alpha\beta}(\gamma^aC)_{\gamma\delta}\Big),\end{aligned}\tag{15}$$

where L_1, L_2 have been found in (12) and L_3 can be obtained as

$$L_3 = (2)^{-2(t+s+u)-1}\pi\frac{\Gamma(-u+\frac{1}{2})\Gamma(-s+\frac{1}{2})\Gamma(-t+\frac{1}{2})\Gamma(-t-s-u)}{\Gamma(-u-t+1)\Gamma(-t-s+1)\Gamma(-s-u+1)}\tag{16}$$

The first terms of $\mathcal{A}_{21}$,$\mathcal{A}_{22}$,$\mathcal{A}_{25}$,$\mathcal{A}_{26}$ ($\mathcal{A}_{23}$) have just infinite t-channel (s-channel) fermion pole and in particular all the terms including the coefficients of L_2 are just contact terms for which they do not have any contribution to the singularities (infinite contact interactions of one RR, two fermions and one scalar field). On the other hand the first term of $\mathcal{A}_{24}$ has just infinite u-channel either gauge or scalar pole.

Notice that the second terms of $\mathcal{A}_{21}$,$\mathcal{A}_{25}$,$\mathcal{A}_{26}$ do involve just infinite s-channel fermion poles as well. Finally the third and fourth terms of $\mathcal{A}_{21}$,$\mathcal{A}_{26}$ are indeed an infinite number of massless $(t+s+u)$-channel poles that in field theory we would clarify which kinds of poles are allowed. The important fact is that these infinite poles should be produced by just either infinite higher derivative corrections to two scalars-two fermions or one scalar/one gauge field and two fermions in IIB. Note that amplitude is antisymmetric with respect to the interchange of fermions as it has already been expected.

Let us make some comments on T-duality. The complete form of this S-matrix can not be obtained by setting T-duality to the previous results (see[22, 23]), since here we do have some terms carrying p^i(closed string momentum in transverse direction), these terms can not derived by applying T-duality , given the fact that RR does not have winding modes in its vertex operator, likewise it is shown in [22] that $C\phi AA$ can not be fully derived from $CAAA$. Thus instead of it we worked with direct computations to find out some special patterns for super string amplitudes.

The expansion is low energy expansion that reflects the fact that all Mandelstam variables should tend to zero $(t, s, u \to 0)$ such that this relation $t+s+u=-p^ap_a$ holds. For further details on the expansions see [12]. The expansions can be restored in some closed form as

$$\begin{aligned}
stL_1 &= -\pi^{3/2}\Big[\sum_{n=-1}^{\infty} b_n\Big(\frac{1}{u}(t+s)^{n+1}\Big) + \sum_{p,n,m=0}^{\infty} e_{p,n,m}u^p(st)^n(s+t)^m\Big] \\
L_3 &= -\frac{\pi^{5/2}}{2}\Bigg(\sum_{n=0}^{\infty} c_n(s+t+u)^n + \frac{\sum_{n,m=0}^{\infty} c_{n,m}[s^nt^m+s^mt^n]}{(t+s+u)} \\
&\quad + \sum_{p,n,m=0}^{\infty} f_{p,n,m}(s+t+u)^p[(s+t)^n(st)^m]\Bigg). \qquad (17)
\end{aligned}$$

Where suL_1 and utL_1can be derived by just replacing $t \leftrightarrow u$ and $s \leftrightarrow u$ within stL_1 expansion and $L_2 = -4(t+s+u)L_3$ which has just contact terms. Some of the coefficients are

$$\begin{aligned}
&b_{-1}=1,\, b_0=0,\, b_1=\frac{1}{6}\pi^2,\, b_2=2\zeta(3), c_0=0, c_1=-\frac{\pi^2}{6}, \\
&e_{2,0,0}=e_{0,1,0}=2\zeta(3), e_{1,0,0}=\frac{1}{6}\pi^2, e_{1,0,2}=\frac{19}{60}\pi^4, e_{1,0,1}=e_{0,0,2}=6\zeta(3), \\
&e_{0,0,1}=\frac{1}{3}\pi^2, e_{3,0,0}=\frac{19}{360}\pi^4, e_{0,0,3}=e_{2,0,1}=\frac{19}{90}\pi^4, e_{1,1,0}=e_{0,1,1}=\frac{1}{30}\pi^4, \qquad (18) \\
&c_2=-2\zeta(3),\, c_{1,1}=\frac{\pi^2}{6},\, c_{0,0}=\frac{1}{2}, f_{0,1,0}=\frac{\pi^2}{3}, f_{0,0,1}=-2\zeta(3)
\end{aligned}$$

It is important to mention that the structure for the coefficients of b_n are the same as the amplitude of one RR and three scalars (three gauge fields) and it has quite universal behaviour[12], however some the coefficients of $c_n, f_{p,n,m}$ do include differences from non-BPS amplitude coefficients [26]. Let us move to field theory section and produce all the infinite u, t, s gauge,scalar/fermion poles. In particular we study different $(u+t+s)-$ channel scalar and gauge poles in order to observe whether or not we can produce all infinite higher derivative corrections of two fermions-two scalars (two fermions-one gauge-one scalar).

3 Infinite scalar $u-$channel poles for $p+2=n$ case

One can expand the closed form of $< V_C V_\phi V_{\bar\psi} V_\psi >$ to actually produce both infinite u-channel (scalar and gauge) poles. It was shown in [12, 14, 22] that the kinetic term of gauge field , tachyons and scalar fields do not get any correction.

Having set the expansions, let us collect all the infinite u-channel scalar poles in string theory as :

$$\begin{aligned}\mathcal{A}_1^{C\phi\bar{\psi}\psi} \sim\; & \frac{ip^i\pi^{3/2}}{2}\sum_{n=-1}^{\infty} b_n\bigg(\frac{1}{u}(t+s)^{n+1}\bigg)(P_-\not{H}_{(n)}M_p)^{\gamma\delta}\xi_{1i}\bar{u}_1^A u_2^B \\ & \times(\gamma^\mu C)_{\gamma\delta}(\gamma_\mu C)_{AB}\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3),\end{aligned} \tag{19}$$

In the above amplitude μ can be both world volume and transverse direction . First let us set it to transverse direction $\mu = j$ and extract the trace as

$$(P_-\not{H}_{(n)}M_p)^{\gamma\delta}(\gamma^j C)_{\gamma\delta} \;=\; \frac{32}{2(p+1)!}(\varepsilon^v)^{a_0\cdots a_p}H^j_{a_0\cdots a_p} \tag{20}$$

Thus replacing the trace in the amplitude we get:

$$\mathcal{A}_1^{C\phi\bar{\psi}\psi} \sim \frac{8ip^i\pi^{3/2}}{(p+1)!}\sum_{n=-1}^{\infty} b_n\bigg(\frac{1}{u}(t+s)^{n+1}\bigg)\xi_{1i}\bar{u}_1^A(\gamma_j)_{AB}u_2^B(\varepsilon^v)^{a_0\cdots a_p}H^j_{a_0\cdots a_p}\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3), \tag{21}$$

One might realize the fact that the same idea (kinetic terms will not receive any correction) must be applied to actually produce all infinite gauge/scalar poles. In the other words in below we will show that the kinetic term of fermion fields has to be fixed and it has no correction at all. The massless pole should be reproduced by the following non-abelian kinetic terms:

$$-T_p\frac{(2\pi\alpha')}{2}\mathrm{Tr}\,\bigg((2\pi\alpha')D_a\phi^i D^a\phi_i + D_aTD^aT - \frac{(2\pi\alpha')}{2}F_{ab}F^{ba} - \bar{\Psi}\gamma^a D_a\Psi\bigg) \tag{22}$$

These infinite u-channel scalar poles can be plotted as follows:



(b)

Figure 1 : The figures related to all infinite u-channel scalar poles of $C\phi\bar{\psi}\psi$

For the field theory of an amplitude including massless scalar fields one must consider three different approaches to end up with their vertices. Basically these methods are either Wess-Zumino (WZ) terms, were introduced by Myers [8], or the needed pull-back methods or the so called Taylor-expansion (given section 5 of [14]). One has to take into account the following field theory amplitude to indeed produce all infinite scalar u-channel poles

$$\mathcal{A} = V^i_\alpha(C_{p+1},\phi_1,\phi)G^{ij}_{\alpha\beta}(\phi)V^j_\beta(\phi,\bar\Psi_1,\Psi_2), \tag{23}$$

Here we have to employ Taylor expansion to obtain the vertex of two scalars and one RR-$p+1$ form field as follows:

$$i\frac{\lambda^2\mu_p}{2!(p+1)!}\int d^{p+1}\sigma(\varepsilon^v)^{a_0\cdots a_p}\,\mathrm{Tr}\,\left(\partial_i\partial_j C^{(p+1)}_{a_0\cdots a_p}\phi^i\phi^j\right) \tag{24}$$

where $\lambda=2\pi\alpha'$, such that we can construct the vertex of one RR,two scalars as

$$V^i_\alpha(C_{p+1},\phi_1,\phi) = i\frac{\lambda^2\mu_p}{2!(p+1)!}(-ip^i)H^j_{a_0\cdots a_p}\xi_{1j}(\varepsilon^v)^{a_0\cdots a_p}\,\mathrm{Tr}\,(\lambda_1\lambda^\alpha) \tag{25}$$

In order to talk about the first simple scalar u-channel pole, one has to take into account the kinetic terms of the scalar and fermion field (the first and last terms in (22)) as well as extracting the covariant derivative of fermion field (as we have just worked out with it in the last section). Hence by considering all possible orderings of scalar and fermions we get it as well as scalar propagator

$$\begin{aligned}V^\beta_j(\bar\Psi_1,\Psi_2,\phi) &= T_p(2\pi\alpha')\bar u^A_1\gamma^j_{AB}u^B_2\left(\mathrm{Tr}\,(\lambda_2\lambda_3\lambda^\beta)-\mathrm{Tr}\,(\lambda_3\lambda_2\lambda^\beta)\right)\\ G^{ij}_{\alpha\beta}(\phi) &= \frac{-i\delta_{\alpha\beta}\delta^{ij}}{T_p(2\pi\alpha')^2k^2}=\frac{-i\delta_{\alpha\beta}\delta^{ij}}{T_p(2\pi\alpha')^2u}\end{aligned} \tag{26}$$

Now if we renormalise the string amplitude(19) by a normalisation factor of $\frac{\mu_p\pi^{-1/2}}{4}$ and replace the above vertices in the field theory amplitude of (23) we observe that, the first simple u-channel pole of string theory ($n=-1$ in string side) can be precisely produced. However notice that the amplitude has infinite u-channel pole now we come to the key point as follows.

The kinetic term of fermion fields has no correction and scalar propagator also does not receive any correction thus the only way to produce all infinite massless poles is to devote infinite higher derivative corrections to the vertex of one RR $p+1$ form field and two scalar fields as follows:

$$i\frac{\lambda^2\mu_p}{2!(p+1)!}\int d^{p+1}\sigma(\varepsilon^v)^{a_0\cdots a_p}\sum_{n=-1}^{\infty}b_n(\alpha')^{n+1}\mathrm{Tr}\,\left(\partial_i\partial_j C^{(p+1)}_{a_0\cdots a_p}D^{a_0}\cdots D^{a_n}\phi^i D_{a_0}\cdots D_{a_n}\phi^j\right) \quad (27)$$

where all the commutators in the covariant derivative of scalar fields should be ignored. Now we can extract the needed vertex as :

$$V^i_\alpha(C_{p+1},\phi_1,\phi)=i\frac{\lambda^2\mu_p}{2!(p+1)!}(-ip^i)H^j_{a_0\cdots a_p}\xi_{1j}(\varepsilon^v)^{a_0\cdots a_p}\sum_{n=-1}^{\infty}b_n(\alpha' k_1.k)^{n+1}\mathrm{Tr}\,(\lambda_1\lambda^\alpha) \quad (28)$$

Having substituted (28) into (23) we get all the infinite scalar u-channel poles as

$$\frac{\alpha'\pi p^i}{(p+1)!}\xi_{1i}\bar{u}^A_1(\gamma_j)_{AB}u^B_2\sum_{n=-1}^{\infty}b_n\frac{(t+s)^{n+1}}{u}(\varepsilon^v)^{a_0\cdots a_p}H^j_{a_0\cdots a_p}\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3), \quad (29)$$

which are all infinite u-channel scalar poles inside the string amplitude as we expected. Hence to provide all infinite scalar poles we had to generalize the vertex of one RR $p+1$-form field and we saw that closed string Ramond-Ramond has induced all infinite higher derivative corrections to two scalar fields. This phenomenon seemed to be universal as it so happens for the other BPS and even non-BPS amplitude (see [12, 22, 23, 14]). In the next section we will see that by applying the same idea and by generalizing all infinite higher derivative corrections of one RR $p-1$ form field ,one gauge and one scalar we could actually produce all infinite u-channel gauge poles of the amplitude as well.

4 Infinite $u-$channel gauge poles for $p=n$ case

By extracting the trace, replacing the desired expansion and setting μ to be located in world volume direction ($\mu=b$) in $\mathcal{A}_{24}$, one can get all the infinite u-channel gauge poles in string theory as follows:

$$\begin{aligned}\mathcal{A}_{24} &\sim \frac{4\mu_p\pi\xi_{1i}(ik_{1a})}{p!}\bar{u}^A_1(\gamma_b)_{AB}u^B_2\sum_{n=-1}^{\infty}b_n\frac{1}{u}(t+s)^{n+1}\\ &\quad\times(\varepsilon^v)^{a_0\cdots a_{p-2}ab}H^i_{a_0\cdots a_{p-2}}\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3)\end{aligned} \quad (30)$$

Note that here we have ignored the second term of $\mathcal{A}_{24}$ which is just contact interaction. The method for deriving contact interactions has been extensively expressed in [12, 14, 22]. These infinite u-channel gauge poles can be drawn as follows :



(b)

Figure 2 : The figures related to all infinite u-channel gauge poles of $C\phi\bar{\psi}\psi$

The following Feynman rule in field theory amplitude for all infinite u-channel gauge poles is needed as

$$\mathcal{A} \;=\; V^a_\alpha(C_{p-1},\phi_1,A)G^{ab}_{\alpha\beta}(A)V^b_\beta(A,\bar{\Psi}_1,\Psi_2), \tag{31}$$

$V^a_\alpha(C_{p-1},\phi_1,A)$ can be derived by making use of the combination of Taylor expansion and WZ terms as

$$i\frac{\lambda^2\mu_p}{p!}\int d^{p+1}\sigma \mathrm{Tr}\left(\partial_i C_{(p-1)}\wedge F\phi^i\right) \tag{32}$$

Therefore one can feasibly derive the following vertex of one RR $p-1$-form field and one off-shell gauge field and one on-shell scalar field as

$$V^a_\alpha(C_{p-1},\phi_1,A) \;=\; i\frac{\lambda^2\mu_p}{p!}H^i_{a_0\cdots a_{p-2}}\xi_{1i}k_{a_{p-1}}(\varepsilon^v)^{a_0\cdots a_{p-1}a}\,\mathrm{Tr}\,(\lambda_1\lambda^\alpha) \tag{33}$$

Now by extracting the kinetic term of fermions, having reminded the covariant derivative of fermion fields ($D^a\psi=\partial^a\psi-i[A^a,\psi]$) and essentially by taking into account all possible orderings of gauge and fermions, we obtain $V^b_\beta(A,\bar{\Psi}_1,\Psi_2)$ as well as gauge propagator

$$\begin{aligned}
V^b_\beta(\bar{\Psi}_1\Psi_2A) &= T_p(2\pi\alpha')\bar{u}^A_1\gamma^b_{AB}u^B_2\left(\mathrm{Tr}\,(\lambda_2\lambda_3\lambda^\beta)-\mathrm{Tr}\,(\lambda_3\lambda_2\lambda^\beta)\right)\\
G^{ab}_{\alpha\beta}(A) &= \frac{-i\delta_{\alpha\beta}\delta^{ab}}{T_p(2\pi\alpha')^2k^2}=\frac{-i\delta_{\alpha\beta}\delta^{ab}}{T_p(2\pi\alpha')^2u}
\end{aligned} \tag{34}$$

with k becomes off-shell gauge field's propagator ($k^a = (k_2 + k_3)^a = (-p - k_1)^a$). Once more by replacing the above vertices in the field theory amplitude of (31) we observe that, the first simple u-channel gauge pole of string theory ($n = -1$ in string side) can be precisely produced, we also used the fact that $p^a(\varepsilon^v)^{a_0\cdots a_{p-1}a} = 0$. However it is worth trying to think of all infinite u-channel poles. We have made some points in the previous section and understood that the kinetic term of fermion fields has no correction and simple scalar/gauge propagators also do not gain any correction, so the only way to produce all infinite u- channel gauge poles is to propose infinite higher derivative corrections to the vertex of one RR $(p-1)-$ form field and one on-shell scalar and an off-shell gauge field as below:

$$i\frac{\lambda^2\mu_p}{p!}\int d^{p+1}\sigma \sum_{n=-1}^{\infty} b_n(\alpha')^{n+1}\mathrm{Tr}\,\left(\partial_i C_{(p-1)}\wedge D^{a_0}\cdots D^{a_n}FD_{a_0}\cdots D_{a_n}\phi^i\right) \tag{35}$$

Now we can extract the needed vertex as :

$$V^a_\alpha(C_{p-1},\phi_1,A) \;=\; i\frac{\lambda^2\mu_p}{p!}H^i_{a_0\cdots a_{p-2}}\xi_{1i}k_{1a_{p-1}}(\varepsilon^v)^{a_0\cdots a_{p-1}a}\sum_{n=-1}^{\infty} b_n(\alpha' k_1.k)^{n+1}\mathrm{Tr}\,(\lambda_1\lambda^\alpha) \tag{36}$$

Having substituted (36) into (31) we get all the infinite u-channel gauge poles of the amplitude as

$$\frac{\alpha'^2\mu_p\pi\xi_{1i}(ik_{1_{a_{p-1}}})}{p!}\bar{u}^A_1(\gamma_a)_{AB}u^B_2\sum_{n=-1}^{\infty} b_n\frac{1}{u}(t+s)^{n+1}(\varepsilon^v)^{a_0\cdots a_{p-1}a}H^i_{a_0\cdots a_{p-2}}\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3) \tag{37}$$

which is exactly all infinite u-channel gauge poles of string amplitude. Hence closed string Ramond-Ramond $(p-1)$-form field has proposed all infinite higher derivative corrections to a scalar and one gauge field. Again we might wonder about this universal behaviour of singularities which does not have any thing to BPS or non-BPS branes .

5 Infinite t, s-channel fermion poles

For the amplitude of one RR, two fermions and one scalar field we realize that there is no graviton propagator in s,t channels either. The reason is that for these channels there is no coupling between one Ramond vertex operator, one scalar and one graviton as one can not saturate the total super ghost charge for disk amplitude, so the only propagator for these channels is indeed fermionic propagator. If we replace usL_1 expansion in the amplitude and simplify it, by extracting all the traces one gets all the infinite t,s-channel fermion poles of the string amplitude as:

$$\begin{aligned}\mathcal{A} &\sim \frac{\alpha'\mu_p\pi\xi_{1i}(2ik_{1a})}{(p+1)!}\bar{u}_1^A(\gamma^a)_{AB}u_2^B\sum_{n=-1}^{\infty}b_n\Big[\frac{1}{t}(u+s)^{n+1}-\frac{1}{s}(u+t)^{n+1}\Big]\\ &\quad\times(\varepsilon^v)^{a_0\cdots a_p}H^i_{a_0\cdots a_p}\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3)\end{aligned}\tag{38}$$

Notice that since the amplitude is totally anti symmetric under interchange of fermionic strings ($s\leftrightarrow t$), there is no need to produce all s-channel poles rather than we just derive all fermionic t-channel poles and finally by interchanging $2\leftrightarrow 3$ and $s\leftrightarrow t$ in all kinematics one can derive all infinite s-channel poles as well. It is worth reminding the point that we have just kept all the singular terms. One can show all the infinite t-channel fermion poles as



(b)

Figure 3 : The figures corresponding to all infinite t-channel fermion poles of $C\phi\bar\psi\psi$

Let us write down the rule to derive all the fermionic t-channel poles

$$\mathcal{A} = V_\alpha(C_{p+1},\Psi_3,\bar\Psi)G_{\alpha\beta}(\Psi)V_\beta(\Psi,\bar\Psi_2,\phi_1),\tag{39}$$

Now by extracting the covariant derivative of fermions in their kinetic term ($D^i\psi=\partial^i\psi-i[\phi^i,\psi]$) and counting all possible orderings of scalar and fermions we get $V_\beta(\Psi,\bar\Psi_2,\phi_1)$ as well as fermion propagator

$$\begin{aligned}V_\beta(\Psi,\bar\Psi_2,\phi_1) &= T_p(2\pi\alpha')\bar{u}_1^A\gamma^j_A\xi_{1j}\Big(\mathrm{Tr}\,(\lambda_2\lambda_2\lambda^\beta)-\mathrm{Tr}\,(\lambda_2\lambda_1\lambda^\beta)\Big)\\ G_{\alpha\beta}(\psi) &= \frac{-i\delta_{\alpha\beta}}{T_p(2\pi\alpha')\not{k}}=\frac{-i\delta_{\alpha\beta}\gamma^a(k_1+k_2)_a}{T_p(2\pi\alpha')t}\end{aligned}\tag{40}$$

Now in order to find $V_\alpha(C_{p+1},\bar\Psi,\Psi)$, one has to find out the new coupling between one RR-$p+1$ form field and two fermions in the world volume of BPS brane which is in fact supersymmtrized version of the known bosonic couplings as follows

$$i\frac{(2\pi\alpha')\mu_p}{(p+1)!}\int d^{p+1}\sigma \mathrm{Tr}\left(C_{a_0\cdots a_p}\bar\Psi\gamma^j\partial_j\Psi\right)(\varepsilon^v)^{a_0\cdots a_p} \tag{41}$$

where equations of motion for fermions must be taken into account ($\not{k}_{2a}\bar{u}=\not{k}_{3a}u=0$). By applying the known field theory techniques the vertex of one RR $p+1$-form field and one off-shell fermion field and one on-shell fermion field can be looked for as follows

$$V_\alpha(C_{p+1},\bar\Psi,\Psi_2) \;=\; i\frac{(2\pi\alpha')\mu_p}{(p+1)!}(\varepsilon^v)^{a_0\cdots a_p}H^i_{a_0\cdots a_p}\gamma^i u_2(\varepsilon^v)^{a_0\cdots a_p}\,\mathrm{Tr}\,(\lambda_3\lambda^\alpha) \tag{42}$$

If we replace above vertices within the field theory amplitude of (39) we observe that, the first simple t-channel pole of string theory ($n=-1$ in string side) shall be resulted. Again in order to actually derive either all infinite $t-$ or s-channel poles, one has to impose the infinite higher derivative interactions to new coupling (41) as well. Thus keeping fixed the kinetic term of fermion field (including no corrections), one has to look for the infinite higher derivative corrections to the vertex of $V_\alpha(C_{p+1},\bar\Psi,\Psi_2)$ as follows

$$i\frac{(2\pi\alpha')\mu_p}{(p+1)!}\int d^{p+1}\sigma\sum_{n=-1}^{\infty}b_n(\alpha')^{n+1}\mathrm{Tr}\left(C_{a_0\cdots a_p}D^{a_0}\cdots D^{a_n}\bar\Psi\gamma^i D_{a_0}\cdots D_{a_n}\partial_i\Psi\right)(\varepsilon^v)^{a_0\cdots a_p} \tag{43}$$

Now by applying the above higher derivative corrections into (39), we are ready to obtain all the infinite t-channel poles. Let us write down the complete form of the $V_\alpha(C_{p+1},\bar\Psi,\Psi_2)$ to all orders in α' as follows:

$$V_\alpha(C_{p+1},\bar\Psi,\Psi) \;=\; i\frac{(2\pi\alpha')\mu_p}{(p+1)!}H^i_{a_0\cdots a_p}\gamma^i u_2(\varepsilon^v)^{a_0\cdots a_p}\,\mathrm{Tr}\,(\lambda_3\lambda^\alpha)\sum_{n=-1}^{\infty}b_n(\alpha' k_3.k)^{n+1} \tag{44}$$

One has to point out the fact the commutator inside the covariant derivative of fermion field should be dropped out. Note also that in producing all the infinite poles in field theory side one has to employ the equations of motion for fermion fields $\not{k}_2\bar{u}=\not{k}_3u=0$ as well.

Setting (44) into (39) one can show that the all the infinite fermion t-channel (s-channel) poles are reconstructed as

$$\begin{aligned}\mathcal{A} &\sim \frac{\alpha'\mu_p\pi\xi_{1i}(2ik_{1a})}{(p+1)!}\bar{u}_1^A(\gamma^a)_{AB}u_2^B\sum_{n=-1}^{\infty}b_n\frac{1}{t}(u+s)^{n+1}\\ &\times(\varepsilon^v)^{a_0\cdots a_p}H^i_{a_0\cdots a_p}\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3)\end{aligned}\tag{45}$$

Therefore the closed string Ramond-Ramond $p+1$-form field has induced an infinite number of higher derivative corrections to two fermions as well as two scalar fields. This clearly confirms that this phenomenon (producing all infinite poles by keeping fixed the kinetic term of open strings and by postulating infinite higher derivative corrections to their vertex operators) is quite universal and might be useful for deriving all the singularities of higher point functions of either BPS or non-BPS amplitudes.

5.1 Infinite higher derivative corrections to two fermions and two scalars and all scalar poles for $p = n - 2$ case

First of all let us write down all the infinite massless scalar $(s+t+u)$-channel poles of the string amplitude of $C\bar{\Psi}\Psi\phi$ for $p = n-2$ case. The goal of this paper is to produce poles order by order and to actually observe whether or not the universal conjecture about higher derivative corrections to all orders in α' (has been made for bosonic amplitude in [23]) can be held for fermionic amplitude? Another goal would be determining all the infinite higher derivative corrections of two fermion two scalars in IIB string theory. Having extracted the traces, one can get to the singularities

$$\begin{aligned}\mathcal{A} &\sim -\frac{\alpha'\pi^{-1/2}\mu_p}{(p+1)!}(\varepsilon^v)^{a_0\cdots a_p}H^i_{a_0\cdots a_p}\xi_{1i}(2ik_{1a})\bar{u}_1^A(\gamma^a)_{AB}u_2^B\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3)\\ &\times\Big[(-t+s)L_3\Big],\end{aligned}\tag{46}$$

The following Feynman diagram must be taken into account



(b)

Figure 4 : The figures corresponding to all infinite $(t+s+u)$-channel scalar poles of $C\phi\bar{\psi}\psi$

The L_3 expansion has infinite $(t+s+u)$- channel pole, let us see its expansion before simplification:

$$L_3 = -\pi^{5/2}\Big(\frac{1}{2(t+s+u)}+\frac{\pi^2(t^2+s^2+u^2)}{12(t+s+u)}+\frac{\xi(3)(t^3+s^3+u^3+tsu)}{(t+s+u)}+\cdots\Big). \quad (47)$$

however, first let us work out just with the first $t+s+u$- channel pole. Reminding the following coupling is worth while for the future field theory amplitude arguments

$$\lambda\mu_p\int d^{p+1}\sigma\frac{1}{(p+1)!}(\varepsilon^v)^{a_0\cdots a_p}\,\mathrm{Tr}\,\left(\phi^i\right)\,H^{(p+2)}_{ia_0\cdots a_p} \quad (48)$$

It is seen in the other amplitudes that the vertex of one RR $p+1$-form field and one scalar field does not get any correction so to produce all infinite $t+s+u$-channel scalar poles, one has to discover the all infinite higher derivative corrections of two fermion and two scalar couplings to all orders in α', in particular in IIB super string theory and in the world volume of just BPS branes. The Feynman rule is given as

$$\mathcal{A} = V^i_\alpha(C_{p+1},\phi)G^{ij}_{\alpha\beta}(\phi)V^j_\beta(\phi,\bar{\Psi},\Psi,\phi_1) \quad (49)$$

where the scalar propagator can be readily derived from the kinetic term of scalar field and $V^i_\alpha(C_{p+1},\phi)$ is derived from the above coupling as

$$\begin{aligned} G^{ij}_{\alpha\beta}(\phi) &= \frac{-i\delta_{\alpha\beta}\delta^{ij}}{T_p(2\pi\alpha')^2k^2}=\frac{-i\delta_{\alpha\beta}\delta^{ij}}{T_p(2\pi\alpha')^2(t+s+u)}, \\ V^i_\alpha(C_{p+1},\phi) &= i(2\pi\alpha')\mu_p\frac{1}{(p+1)!}(\varepsilon^v)^{a_0\cdots a_p}H^i_{a_0\cdots a_p}\mathrm{Tr}\,(\lambda_\alpha). \end{aligned} \quad (50)$$

Notice that scalar in $V^i_\alpha(C_{p+1},\phi)$ must be abelian so $\mathrm{Tr}\,(\lambda_\alpha)$ makes sense for the abelian matrix λ_α.

By taking a look at (46) one understands that the first simple $(t+s+u)$-scalar pole has to be discovered by a coupling between two scalars and two fermions such that they should carry three momenta. One can see that the kinematic factor of $(-t+s)$ get factorized and this is in favor of the given universal conjecture for all infinite higher derivative corrections [23]. First we write down the most possible desired couplings as follow:

$$\frac{T_p(2\pi\alpha')^3}{4}(\bar{\Psi}\gamma^aD_b\Psi D^a\phi^iD^b\phi_i+D^a\phi^iD^b\phi_i\bar{\Psi}\gamma^aD_b\Psi), \qquad \frac{T_p(2\pi\alpha')^3}{8}\bar{\Psi}\gamma^aD_a\Psi D^b\phi^iD_b\phi_i \quad (51)$$

The third coupling is gotten after the multiplication of the kinetic terms of fermions and scalars and using the desired expansions (their coefficients were fixed by comparing them with our S-matrix (46), for more details see [23]). Note that all the connections(commutator terms in the definition of the covariant derivative of fermion fields must be overlooked). Let us work out with $(\bar{\Psi}\gamma^a D_b\Psi D^a\phi^i D^b\phi_i + D^a\phi^i D^b\phi_i\bar{\Psi}\gamma^a D_b\Psi)$. In order to produce the field theory vertex operators , one has to consider $\mathrm{Tr}\,(\lambda_2\lambda_3\lambda_1\lambda_\beta), \mathrm{Tr}\,(\lambda_2\lambda_3\lambda_\beta\lambda_1)$ for the first coupling and two different orderings for the second coupling as below: four different possible orderings as

$$\mathrm{Tr}\,(\lambda_\beta\lambda_1\lambda_2\lambda_3), \mathrm{Tr}\,(\lambda_1\lambda_\beta\lambda_2\lambda_3) \tag{52}$$

where λ_β is related to Abelian scalar field in the propagator. By extracting the first and second coupling in (51) and considering all orderings one obtains

$$V^j_\beta(\phi,\bar{\Psi},\Psi,\phi_1) = i\frac{T_p(2\pi\alpha')^3}{4}\bar{u}^A(\gamma^a)_{AB}u^B\xi_{1j}\Big(-k_{1a}\frac{t}{2}-k_{4a}\frac{s}{2}\Big)\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3\lambda_\beta)$$

Now one has to apply momentum conservation along the world volume of brane and makes use of the equations of motion for fermion fields such that at the end we are left with the following result :

$$V^j_\beta(\phi,\bar{\Psi},\Psi,\phi_1) = i\frac{T_p(2\pi\alpha')^3}{4}k_{1a}\bar{u}^A(\gamma^a)_{AB}u^B\xi_{1j}(-\frac{t}{2}+\frac{s}{2})\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3\lambda_\beta) \tag{53}$$

Now if we replace (53) into (49) and also substitute (47) inside (46) , we will observe that exactly the first (t+s+u)-channel scalar pole in (46) is produced. However as it is clear from the L_3 expansion , it has infinite massless (t+s+u)-channel scalar poles. Keeping in mind the fact that $V^i_\alpha(C_{p+1},\phi)$ and the simple scalar propagator do not require any corrections , one would expect that all infinite poles are related to the infinite higher derivative corrections of the two fermions and two scalar fields in IIB Super string theory. Let us move to the second coupling. For $\bar{\Psi}\gamma^a D_a\Psi D^b\phi^i D_b\phi_i$ coupling one has to once more consider two different possible orderings as

$$\mathrm{Tr}\,(\lambda_2\lambda_3\lambda_1\lambda_\beta), \qquad \mathrm{Tr}\,(\lambda_2\lambda_3\lambda_\beta\lambda_1) \tag{54}$$

where λ_β is related to Abelian scalar field in the propagator. By extracting this coupling and considering all orderings one gets

$$V^j_\beta(\phi,\bar{\Psi},\Psi,\phi_1) = -ik_{3a}\bar{u}^A(\gamma^a)_{AB}u^B\xi_{1j}u\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3\lambda_\beta)$$

one may apply on-shell condition $(t+s+u=0)$, however this coupling is vanished by applying the equation of motion for fermion field ($\not{k}_3 u = 0$) so it does not have any contribution to field theory amplitude. Let us now generalize our method to find out all infinite couplings (to all orders in α') to produce all the infinite $(t+s+u)$-channel scalar poles. Having taken the following couplings

$$\mathcal{L}^{n,m} = \pi^3\alpha'^{n+m+3}T_p\Big(a_{n,m}\mathrm{Tr}\Big[\mathcal{D}_{nm}\left(\bar\Psi\gamma^a D_b\Psi D^a\phi^i D^b\phi_i\right) + \mathcal{D}_{nm}\left(D^a\phi^i D^b\phi_i\bar\Psi\gamma^a D_b\Psi\right)$$
$$+h.c\Big] + ib_{n,m}\mathrm{Tr}\Big[\mathcal{D}'_{nm}\left(\bar\Psi\gamma^a D_b\Psi D^a\phi^i D^b\phi_i\right) + \mathcal{D}'_{nm}\left(D^a\phi^i D^b\phi_i\bar\Psi\gamma^a D_b\Psi\right) + h.c.\Big]\Big) \quad (55)$$

with the definition of higher derivative operator of $\mathcal{D}_{nm}$ as

$$\begin{aligned}\mathcal{D}_{nm}(EFGH) &\equiv D_{b_1}\cdots D_{b_m}D_{a_1}\cdots D_{a_n}EFD^{a_1}\cdots D^{a_n}GD^{b_1}\cdots D^{b_m}H\\ \mathcal{D}'_{nm}(EFGH) &\equiv D_{b_1}\cdots D_{b_m}D_{a_1}\cdots D_{a_n}ED^{a_1}\cdots D^{a_n}FGD^{b_1}\cdots D^{b_m}H\end{aligned}$$

we now show that one can precisely produce all infinite scalar poles. Let us focus on $a_{n,m}$. Applying standard field theory techniques, considering hermitian conjugate of the above couplings, concerning momentum conservation in world volume direction, using the equations of motion for fermion fields and finally taking into account all possible different orderings[3] one obtains

$$V^j_\beta(\phi,\bar\Psi,\Psi,\phi_1) = iT_p\frac{(2\pi\alpha')^3}{4}k_{1a}\bar u^A(\gamma^a)_{AB}u^B\xi_{1j}\Big(-\frac{t}{2}t^n s^m + \frac{s}{2}t^m s^n\Big)\mathrm{Tr}\left(\lambda_1\lambda_2\lambda_3\lambda_\beta\right) \quad (56)$$

Now if we replace (56) into (49) , we will observe that exactly all the infinite (t+s+u)-channel scalar poles in (46) can be produced. For instance by putting $n,m=0$ we have already in this section shown that the first $(t+s+u)$ channel scalar pole has been produced. Let us move on with $n=1, m=0$ case. From field theory side we get

$$V^j_\beta(\phi,\bar\Psi,\Psi,\phi_1) = iT_p\frac{(2\pi\alpha')^3}{4}k_{1a}\bar u^A(\gamma^a)_{AB}u^B\xi_{1j}\Big(-\frac{t^2}{2} + \frac{s^2}{2}\Big) \quad (57)$$

However note that again the over all coefficient of string amplitude $(k_{1a}(s-t))$ can be extracted from the above vertex and the the rest will be produced by taking the coefficient of $c_{1,1}$ in string amplitude. Note that if we concentrate on $b_{n,m}$ and extract the needed vertex we will get

$$V^j_\beta(\phi,\bar\Psi,\Psi,\phi_1) = iT_p\frac{(2\pi\alpha')^3}{4}k_{1a}\bar u^A(\gamma^a)_{AB}u^B\xi_{1j}\Big(-\frac{t}{2}u^n s^m + \frac{s}{2}t^m u^n\Big)\mathrm{Tr}\left(\lambda_1\lambda_2\lambda_3\lambda_\beta\right) \quad (58)$$

---

[3] Notice the fact that BPS branes do not carry Chan -Paton factor so we do not expect to have any $(-\mathcal{D}_{nm}), (-\mathcal{D}'_{nm})$ operators in the above couplings(the same facts were found for bosonic amplitudes (see [12])).

However one has to apply on-shell condition $s+t+u=0$ into above vertex to be able to produce all infinite poles. Indeed similar checks for higher order α' corrections and in particular for the other amplitudes in [12, 22] have been carried out. This ends our goal for obtaining an infinite number of higher derivative corrections of two fermions and two scalars to all orders in α'.

5.2 The higher derivative corrections to two fermions, one gauge and one scalar field and the infinite gauge poles for $p=n$ case

The final sinularities in the amplitude of $C\bar\Psi\Psi\phi$ are related to all infinite massless $(s+t+u)$-channel gauge poles of the string theory for $p=n$ case. The goal here is to actually find out the first non zero couplings of one gauge,one scalar and two fermions of IIB string theory and to fix their coefficients by comparing field theory amplitude with string theory amplitude. Of course, essentially one would like to derive infinite corrections which would be nice to obtain them and to see whether or not the universal conjecture about higher derivative corrections to all orders in α' [23] holds for fermionic amplitude? Last aim is related to fixing the coefficients of all needed corrections which must be looked for by comparsion with in string theory amplitude and not by any other tools such as T-duality.

Extracting the traces one can find those terms as

$$\mathcal{A} \sim -\frac{\alpha'\pi^{-1/2}\mu_p}{(p)!}(\varepsilon^v)^{a_0\cdots a_{p-1}a}H_{a_0\cdots a_{p-1}}\xi_{1i}(2ik_{1a})\bar{u}_1^A(\gamma^i)_{AB}u_2^B\,\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3)\bigg[-2tL_3\bigg], \tag{59}$$

The corresponding Feynman diagram can be drawn as



Figure 5 : The figures corresponding to all infinite gauge $(t+s+u)$-channel of $C\phi\bar\psi\psi$

We have already pointed out that the expansion of the L_3 had infinite $(t+s+u)$- channel poles. Let us first carry out field theory argument to produce just the first simple $t+s+u$-gauge pole. The following coupling has been derived in [12] such that

$$i(2\pi\alpha')\mu_p\int d^{p+1}\sigma\frac{1}{(p)!}C_{p-1}\wedge F \tag{60}$$

It has already been discovered in the amplitude of one RR and three gauge fields [12] that the vertex of one RR $(p-1)$-form field and one gauge field $V(C_{p-1},A)$ does not get any correction so one expects that to produce all the infinite gauge poles for $p=n$ case the only corrections are turned to the corrections of one off-shell gauge, one on-shell scalar field and two fermion fields (to all orders in α' in IIB super string theory). Thus the infinite higher derivative corrections should be explored by matching all the infinite gauge poles of $C\bar{\Psi}\Psi\phi$ for $p=n$ case with field theory amplitude.

The Feynman rule is

$$\mathcal{A} \;=\; V^a_\alpha(C_{p-1},A)G^{ab}_{\alpha\beta}(A)V^b_\beta(A,\bar{\Psi},\Psi,\phi_1) \tag{61}$$

where the gauge propagator can be obtained from the kinetic term of gauge field and $V^a_\alpha(C_{p-1},A)$ is gotten from the (60)

$$\begin{aligned}G^{ab}_{\alpha\beta}(A) &= \frac{-i\delta_{\alpha\beta}\delta^{ab}}{T_p(2\pi\alpha')^2k^2}=\frac{-i\delta_{\alpha\beta}\delta^{ab}}{T_p(2\pi\alpha')^2(t+s+u)},\\ V^a_\alpha(C_{p-1},A) &= i(2\pi\alpha')\mu_p\frac{1}{(p)!}(\varepsilon^v)^{a_0\cdots a_{p-1}a}H_{a_0\cdots a_{p-1}}\,\mathrm{Tr}\,(\lambda_\alpha).\end{aligned} \tag{62}$$

Notice that the gauge field in $V^a_\alpha(C_{p-1},A)$ must be abelian so $\mathrm{Tr}\,(\lambda_\alpha)$ makes sense just for the abelian matrix λ_α.

By taking a look at (59) one understands that the first simple $(t+s+u)$-scalar pole has to be discovered by a coupling between a scalar,a gauge and two fermions such that they should carry three momenta. Consider the following couplings:

$$\frac{T_p(2\pi\alpha')^3}{4}\Big[\bar{\Psi}\gamma^iD_b\Psi D^a\phi^iF_{ab}+\bar{\Psi}\gamma^iD_b\Psi F_{ab}D^a\phi^i\Big] \tag{63}$$

which are the multiplication of the kinetic terms of fermions and the field strength of gauge field and the covariant derivative of the scalar field (we have set them to have general covariance). We have to highlight the fact that the connections(commutator terms in the definition of the covariant derivative of fermion fields and in scalar fields must be overlooked). Let us work out with (63). Unlike the previous section, here the only possible orderings for the first and second term of (63) accordingly are

$$\mathrm{Tr}\,(\lambda_2\lambda_3\lambda_1\lambda_\beta), \qquad\qquad \mathrm{Tr}\,(\lambda_2\lambda_3\lambda_\beta\lambda_1)$$

where λ_β is related to Abelian gauge field in the propagator. By extracting these couplings

and applying momentum conservation along the world volume of brane and making use of the equations of motion for fermion fields we obtain :

$$V^b_\beta(A,\bar\Psi,\Psi,\phi_1) \;=\; i\frac{T_p(2\pi\alpha')^3}{2}\bar u^A(\gamma^j)_{AB}u^B\xi_{1j}\Big(k_{1b}\frac{t}{2}\Big)\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3\lambda_\beta) \tag{64}$$

Now if we replace (64) into (61) and also substitute (47) inside (59) , we will observe that exactly the first (t+s+u)-channel gauge pole in (59) is produced. However, as it is clear from the L_3 expansion , it has infinite massless (t+s+u)-channel gauge poles. Keeping in mind the fact that $V^a_\alpha(C_{p-1},A)$ and the simple gauge propagator do not require any corrections , one would expect that all infinite poles are related to the infinite higher derivative corrections of two fermions and one scalar and one gauge field in IIB Super string theory. We have already introduced how to look for higher derivative corrections. One can generalize the couplings in (63) to higher derivative corrections as well. Given the leading couplings in (63), one can discover all order couplings as

$$\begin{aligned}\mathcal{L}^{n,m} &= \pi^3\alpha'^{n+m+3}T_p\Big(a_{n,m}\mathrm{Tr}\Big[\mathcal{D}_{nm}\left(\bar\Psi\gamma^iD_b\Psi D^a\phi^iF_{ab}\right)+\mathcal{D}_{nm}\left(\bar\Psi\gamma^iD_b\Psi F_{ab}D^a\phi^i\right)\\ &+h.c\Big]+ib_{n,m}\mathrm{Tr}\Big[\mathcal{D}'_{nm}\left(\bar\Psi\gamma^iD_b\Psi D^a\phi^iF_{ab}\right)+\mathcal{D}'_{nm}\left(\bar\Psi\gamma^iD_b\Psi F_{ab}D^a\phi^i\right)+h.c.\Big]\Big)\end{aligned}$$

where now for the terms including $a_{n,m}$ coefficients one must replace the following coupling

$$V^b_\beta(A,\bar\Psi,\Psi,\phi) = i\frac{T_p(2\pi\alpha')^3}{4}\bar u^A(\gamma^j)_{AB}u^B\xi_{1j}\Big(k_{1b}\frac{t}{2}\Big)(t^ns^m+s^nt^m)\mathrm{Tr}\,(\lambda_1\lambda_2\lambda_3\lambda_\beta) \tag{65}$$

rather than (64) into (61) to be able to give rise all the infinite gauge poles of the amplitude. Finally it is worth mentioning the fact that for $b_{n,m}$ coefficients one must apply on-shell condition ($t+s+u=0$) at each order of α' to get to the higher derivative corrections. It would be nice to find the higher derivative corrections to two fermions and two gauge fields to all orders in α' [28].

6 Conclusions

By applying direct CFT techniques we found the complete form of the $< V_C V_{\bar{\psi}} V_\psi V_\phi >$ amplitude in IIB. All infinite scalar/gauge(for $p+2=n, p=n$ cases) and fermion poles have been explored. The vertices of $V_i^\alpha(C_{p+1},\phi), V_a^\alpha(C_{p-1},A)$ do not require any corrections so all infinite $(t+s+u)-$ channel scalar/gauge poles have provided worth information to determine infinite higher derivative corrections (to all orders in α') to two fermions and two scalars and also to two fermions, one scalar and one gauge couplings for which we derived them in this paper. We also clarified that the same universal conjecture [23] for all higher derivative corrections holds even for fermionic amplitudes in IIB.

It is worth pointing out that in RR vertex operator there are no winding modes so applying T-duality is not so effective and in particular in order not to miss any terms in the amplitudes and to get to all higher derivative corrections with their exact coefficients one has to apply direct computations and we proposed some patterns instead of it.

Let us talk about a subtle issue regarding the relation of open/closed string vertices in II super string theory. For our amplitude which involves mixed open / closed strings, our calculations make sense just by using path integral formalism such that propagators (Green functions) are found by CFT methods while closed string has $(\alpha_n, \tilde{\alpha}_n)$ oscillators. It is not obvious how to do calculations with oscillators, namely in the first string quantization it is subtle how to deal with $\tilde{\alpha}_n$. Some comments are suggested in (3.4) of [37] such that both oscillators for closed string are determined with open string's ones. In the other words some analytic continuation is needed and this kind of realization implies that the state for closed string as a composite state of the open strings. The interpretation in field theory might be useful mentioning.

It reveals that all background fields in DBI should be some functions of SYM fields or in the other words background fields must become composite and finally these functions would be examined once we work out with the complete open string formalism. Note also that super gravity background fields should have included Taylor expansion as was argued in [8].

Acknowledgment

I would like to thank J.Polchinski,J.Maldacena, R.Myers, W.Lerche,S.Stieberger,G.Veneziano, K.S.Narain and N.Lambert for having valuable discussions during several conferences.

References

[1] J. Polchinski,"Dirichlet-Branes and Ramond-Ramond Charges," Phys. Rev. Lett. **75**, 4724 (1995) [arXiv:hep-th/9510017].

[2] E. Witten,"Bound states of strings and p-branes," Nucl. Phys. B **460**,335 (1996) [arXiv:hep-th/9510135].

[3] J. Polchinski," Lectures on D-branes," [arXiv:hep-th/9611050] ; C. P. Bachas, "Lectures on D-branes," [arXiv:hep-th/9806199].

[4] J. Polchinski, "String duality: A Colloquium," Rev. Mod. Phys. **68**, 1245 (1996) [hep-th/9607050].

[5] E. Hatefi, A. J. Nurmagambetov and I. Y. Park, "N^3 entropy of $M5$ branes from dielectric effect," Nucl. Phys. B **866**, 58 (2013) [arXiv:1204.2711 [hep-th]].

[6] E. Hatefi, A. J. Nurmagambetov and I. Y. Park, "ADM reduction of IIB on $H^{p,q}$ to dS braneworld," arXiv:1210.3825 [hep-th].

[7] J. Polchinski, "Combinatorics of boundaries in string theory," Phys. Rev. **D50**,6041 (1994)[arXiv:hep-th/9407031] ; J. Polchinski, S. Chaudhuri and C. V. Johnson, "Notes on D-Branes," [arXiv:hep-th/9602052].

[8] R. C. Myers,"Dielectric-branes," JHEP **9912**, 022 (1999) [arXiv:hep-th/9910053].

[9] P. S. Howe, U. Lindstrom and L. Wulff, 'On the covariance of the Dirac-Born-Infeld-Myers action," JHEP **0702**, 070 (2007) [hep-th/0607156].

[10] R. G. Leigh, "Dirac-Born-Infeld Action from Dirichlet Sigma Model," Mod. Phys. Lett. A **4**, 2767 (1989).

[11] M. Cederwall, A. von Gussich, B. E. W. Nilsson and A. Westerberg, "The Dirichlet super three-brane in ten-dimensional type IIB supergravity," Nucl. Phys. B **490**, 163

(1997) [hep-th/9610148] ; M. Aganagic, C. Popescu and J. H. Schwarz, "D-brane actions with local kappa symmetry," Phys. Lett. B **393**, 311 (1997) [hep-th/9610249] ; M. Aganagic, C. Popescu and J. H. Schwarz, "Gauge invariant and gauge fixed D-brane actions," Nucl. Phys. B **495**, 99 (1997) [hep-th/9612080] ; M. Cederwall, A. von Gussich, B. E. W. Nilsson, P. Sundell and A. Westerberg, Nucl. Phys. B **490**, 179 (1997) [hep-th/9611159] ; E. Bergshoeff and P. K. Townsend, Nucl. Phys. B **490** (1997) 145 [hep-th/9611173].

[12] E. Hatefi, "On effective actions of BPS branes and their higher derivative corrections," JHEP **1005**, 080 (2010) [arXiv:1003.0314 [hep-th]].

[13] E. Hatefi, "Shedding light on new Wess-Zumino couplings with their corrections to all orders in alpha-prime," arXiv:1211.2413 [hep-th].

[14] E. Hatefi, "On higher derivative corrections to Wess-Zumino and Tachyonic actions in type II super string theory," Phys. Rev. D **86**, 046003 (2012) [arXiv:1203.1329 [hep-th]].

[15] I. Y. Park, "Scattering on D3-branes," Phys. Lett. **B660**, 583 (2008) [arXiv:0708.3452[hep-th]] ; I. Y. Park,"Open string engineering of D-brane geometry," JHEP **0808**, 026 (2008) [arXiv:0806.3330[hep-th]].

[16] P. Koerber and A. Sevrin, "The NonAbelian D-brane effective action through order alpha-prime**4," JHEP **0210**, 046 (2002) [hep-th/0208044].

[17] A. Keurentjes, P. Koerber, S. Nevens, A. Sevrin and A. Wijns, "Towards an effective action for D-branes," Fortsch. Phys. **53**, 599 (2005) [hep-th/0412271].

[18] F. Denef, A. Sevrin and J. Troost, "NonAbelian Born-Infeld versus string theory," Nucl. Phys. B **581**, 135 (2000) [hep-th/0002180].

[19] A. Hashimoto and I. R. Klebanov, "Scattering of strings from D-branes," Nucl. Phys. Proc. Suppl. **55B**, 118 (1997) [arXiv:hep-th/9611214] ; A. Hashimoto and I. R. Klebanov , "Decay of excited D-branes," Phys. Lett. **B381**, 437 (1996) [arXiv:hep-th/9604065] ; I. R. Klebanov and L. Thorlacius, "The Size of p-branes," Phys. Lett. **B371**, 51 (1996)[arXiv:hep-th/9510200] ; S.S. Gubser, A. Hashimoto, I.R. Klebanov, and J.M. Maldacena, "Gravitational lensing by p-branes,"Nucl. Phys. **B472**, 231 (1996) [arXiv:hep-th/9601057] ; C. Bachas, "D-Brane Dynamics," Phys.

Lett. **B374**, 37 (1996)[arXiv:hep-th/9511043] ; J. Polchinski,“String duality: A colloquium,” Rev. Mod. Phys. **68**, 1245 (1996) [arXiv:hep-th/9607050]; W. Taylor,“Lectures on D-branes, gauge theory and M(atrices),” [arXiv:hep-th/9801182]; C. Vafa,“Lectures on strings and dualities,” [arXiv:hep-th/9702201] ; ;M. Billo, M. Frau, F. Lonegro and A. Lerda,“N = 1/2 quiver gauge theories from open strings with R-R fluxes,” JHEP **0505**,047 (2005) [arXiv:hep-th/0502084] ; M. Billo, P. Di Vecchia, M. Frau, A. Lerda, I. Pesando, R. Russo and S. Sciuto,“Microscopic string analysis of the D0-D8 brane system and dual R-R states,” Nucl. Phys. B **526**,199 (1998) [arXiv:hep-th/9802088]. ; E. Hatefi, “Three Point Tree Level Amplitude in Superstring Theory,” Nucl. Phys. Proc. Suppl. **216**, 234 (2011) [arXiv:1102.5042 [hep-th]].

[20] E. Hatefi, A. J. Nurmagambetov and I. Y. Park, “Near-Extremal Black-Branes with n*3 Entropy Growth,” Int. J. Mod. Phys. A **27**, 1250182 (2012) [arXiv:1204.6303 [hep-th]]

[21] J. McOrist and S. Sethi, “M-theory and Type IIA Flux Compactifications,” JHEP **1212**, 122 (2012) [arXiv:1208.0261 [hep-th]].

[22] E. Hatefi and I. Y. Park, “More on closed string induced higher derivative interactions on D-branes,” Phys. Rev. D **85**, 125039 (2012) [arXiv:1203.5553 [hep-th]].

[23] E. Hatefi and I. Y. Park, “Universality in all-order α' corrections to BPS/non-BPS brane world volume theories,” Nucl. Phys. B **864** (2012) 640 [arXiv:1205.5079 [hep-th]].

[24] E. Hatefi, “On D-brane anti D-brane effective actions and their corrections to all orders in alpha-prime,” arXiv:1211.5538 [hep-th].

[25] M. R. Garousi and E. Hatefi,“On Wess-Zumino terms of Brane-Antibrane systems,” Nucl. Phys. B **800**, 502 (2008) [arXiv:0710.5875 [hep-th]].

[26] M. R. Garousi and E. Hatefi,“More on WZ actions of non-BPS branes,” JHEP **0903**, 08 (2009) [arXiv:0812.4216 [hep-th]].

[27] C. Vafa, “Evidence for F theory,” Nucl. Phys. B **469**, 403 (1996) [hep-th/9602022].

[28] E. Hatefi, “Work in progress”

[29] S. Stieberger,“ Open, Closed vs.Pure Open String Disk Amplitudes,” [arXiv:0907.2211 [hep-th]].

[30] C. Kennedy and A. Wilkins,“Ramond-Ramond couplings on brane-antibrane systems,” Phys. Lett. B **464**, 206 (1999) [arXiv:hep-th/9905195].

[31] L. A. Barreiro and R. Medina,“5-field terms in the open superstring effective action,” JHEP **0503**,055 (2005) [arXiv:hep-th/0503182]. ; R. Medina, F. T. Brandt and F. R. Machado,“The open superstring 5-point amplitude revisited,” JHEP **0207**,071 (2002) [arXiv:hep-th/0208121] ; L. A. Barreiro and R. Medina, “Revisiting the S-matrix approach to the open superstring low energy effective lagrangian,” JHEP **1210**, 108 (2012) [arXiv:1208.6066 [hep-th]] ; A. Bilal, “Higher derivative corrections to the nonAbelian Born-Infeld action,” Nucl. Phys. B **618**, 21 (2001) [hep-th/0106062].

[32] J. Polchinski,“String theory ”,Vol 2,Cambridge University Press,1998

[33] H. Liu and J. Michelson, “*-trek 3: The Search for Ramond-Ramond couplings,” Nucl. Phys. B **614**, 330 (2001) [hep-th/0107172].

[34] V. A. Kostelecky, O. Lechtenfeld, W. Lerche, S. Samuel and S. Watamura, “Conformal Techniques, Bosonization and Tree Level String Amplitudes,” Nucl. Phys. B **288**, 173 (1987).

[35] D. Haertl and O. Schlotterer, “Higher Loop Spin Field Correlators in Various Dimensions,” Nucl. Phys. B **849** (2011) 364 [arXiv:1011.1249 [hep-th]].

[36] A. Fotopoulos,“On (alpha’)**2 corrections to the D-brane action for non-geodesic world-volume embeddings,” JHEP **0109**, 005 (2001) [arXiv:hep-th/0104146].

[37] M. Billo, M. Frau, F. Fucito and A. Lerda, “Instanton calculus in R-R background and the topological string,” JHEP **0611**, 012 (2006) [hep-th/0606013].